



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A second group, among which are the common cereals and various other plants, as *Lactuca sativa*, *Helianthus annuus*, *Asparagus officinalis*, *Pinus virginiana*, *Robinia pseudacacia*, either all decayed before germinating or germinated and then all decayed before being examined. A third group, which includes our more noxious weeds, retained their vitality to a considerable degree. The deeper the seeds were buried the better they retained their vitality. Vitality is best preserved, even in weed seeds, when they are carefully harvested and stored in a dry and comparatively cool place.—WM. CROCKER.

Prothallia and sporelings of *Botrychium*.—BRUCHMANN³⁵ has been investigating *Botrychium Lunaria*. Since this species has no means of vegetative multiplication, like the adventitious shoots of *Ophioglossum vulgatum*, every sporophyte must have come from a gametophyte. The prothallia are hard to find because they are very small (1–2^{mm} long and 0.5–1^{mm} wide), and the sporelings grow for several years before they reach the surface of the soil. The prothallia are found at a depth of 1–3^{cm}. In form and general character the prothallium of *B. Lunaria* resembles that of *B. virginianum*, except that it is much smaller. BRUCHMANN succeeded in germinating the spores and his results agree with those of CAMPBELL, who got the two and three-cell stage in *Ophioglossum vulgatum*. Further work upon this aspect of the problem will be published later. However, he represents a single cell at the “spore pole” of the prothallium and regards this as the first cell of the prothallium, representing the protonema stage. Nearly every prothallium bears an embryo and some prothallia have two. The first division of the embryo is transverse. Growth is very slow, the sporeling being three years old before it reaches the surface. One plate and considerable attention in the text is devoted to the anatomy of the mature plant.—CHARLES J. CHAMBERLAIN.

Spermatozoids of *Cycas revoluta*.—MIYAKE³⁶ studied the living spermatozoids at the island of Oshima (28° 30' N) in September, and in southern Japan (31° 35' N) from the beginning to the middle of October. The diameter of the spermatozoids varies from 180 to 210 μ . The two spermatozoids are surrounded by a delicate membrane, but it could not be determined with certainty whether the membrane belongs to the spermatozoid or is merely the *Hautschicht* of the protoplasm of the pollen tube. For observing the movements the spermatozoids were placed in a 10 per cent. cane sugar solution. The movements often continued for one to three hours; and in one case for six hours and forty minutes, and in another case for five hours and thirty minutes. In some cases the spermatozoids were shot out suddenly from the pollen tube, which seems to be the method that occurs under natural conditions. The forward movement is always accompanied by a rotation from left to right about the

³⁵ BRUCHMANN, H., Ueber das Prothallium und die Sporenpflanze von *Botrychium Lunaria* Sw. Flora 96:203–230. pls. 1–2. 1906.

³⁶ MIYAKE, Ueber die Spermatozoiden von *Cycas revoluta*. Ber. Deutsch. Bot. Gesell. 24:78–83. pl. 6. 1906.

axis. In some cases the forward movement was found to be at the rate of 0.7^{mm} per second. MIYAKE agrees with WEBBER that the liquid in the archegonial chamber at the time of fertilization comes from the pollen tube and not from the archegonium.—CHARLES J. CHAMBERLAIN.

Heterostyly and gynodioecism.—Inheritance of dimorphism has been investigated by RAUNKIÄR³⁷ in *Primula*, *Menyanthes*, *Pulmonaria*, *Fagopyrum*, *Knautia*, and *Thymus*. In all heterostylic species studied he finds that the long-styled and short-styled forms occur in about equal numbers regardless of the character of the environment. In gynodioecious species, on the other hand, he finds considerable variation in the proportions of the two forms in different localities. The results of breeding are in close accord with those of CORRENS,³⁸ except in an interesting case in which a cross between two bisporangiate plants of *Thymus vulgaris* produced 65 per cent. pistillate plants. In *Primula officinalis*, brachystylic plants pollinated by brachystylic produced 62.5 per cent. brachystylic, brachystylic \times dolichostylic gave 55.2 per cent. brachystylic, and dolichostylic \times dolichostylic only 4.3 per cent. brachystylic. Investigation covering several generations is needed to determine the effects of the pre-parental ancestry, and until this is done, any speculation as to the hereditary nature of the forms of a dimorphic species can be of little value.—GEORGE H. SHULL.

Development of spores of Helminthostachys.—BEER³⁹ has investigated the development of the spores of *H. zeylanica*, his material being fertile spikes preserved in spirit. CARDIFF,⁴⁰ and afterwards STEVENS,⁴¹ had described the peculiar blocking out of the sporogenous tissue and the remarkable behavior of the plasmodium-like tapetal cytoplasm in *Botrychium*; and BEER finds the same phenomena in *Helminthostachys*. His observations extend, however, to the specific work of the tapetal plasmodium in spore-formation. The observed facts are that during the period of exospore growth the tapetal plasmodium shows more or less complete disappearance of starch, gradual diminution of the finely vacuolar cytoplasm, and richly chromatic nuclei which often show irregularities of outline. The conclusion is that the tapetal plasmodium is the center of metabolic activities in which a substance is elaborated from the raw materials contained in the tapetum, and is employed, directly or indirectly, in the growth of the spore wall.—J. M. C.

³⁷ RAUNKIÄR, C., Sur la transmission par hérédité dans les espèces hétéromorphes. Bull. Acad. Roy. Sci. et Let., Denmark, pp. 31–39, 1906.

³⁸ See BOT. GAZETTE 39: 304, 1905, and 41: 302, 1906.

³⁹ BEER, RUDOLF, On the development of the spores of *Helminthostachys zeylanica*. Annals of Botany 20:177–186. pls. 11–12. 1906.

⁴⁰ BOT. GAZETTE 29:340–347. pl. 9. 1905.

⁴¹ Annals of Botany 19:—. —. 1905.